

Effect of Urea Fertilizer on Growth and Yield of Sorghum (*Sorghum bicolor* L.) Under ~~Reined~~ Conditions, ~~on~~ North Kordofan, Sudan.

Delete and replace with rain- :[1GG]Comment
fed

Delete and replace with " in " :[2GG]Comment

Comment [GG3]:
Effect of Urea Fertilizer on Growth and Yield components of Sorghum (*Sorghum bicolor* L.) Under ~~rain-fed~~ Conditions, ~~in~~ North Kordofan, Sudan.

ABSTRACT

Low soil fertility is one of the main factors ~~responsible~~ for low productivity of crops in North Kordofan State. The effects of urea fertilizer on growth and yield of sorghum (*Sorghum bicolor* L.) variety *Botana* was studied at the experimental field of Faculty of Natural Resources and Environmental Studies, in Sheikan locality, North Kordofan State during two consecutive rainy seasons 2021/2022 and 2022/2023. The experiment consisted of three treatments (Control, 40 kg urea/feddan and 80 kg urea /feddan). Treatments were laid out in a Randomized Complete Block Design (RCBD) with four replications. Growth characters studied were (days to 50% flowering, plant height (cm), number of leaf per plant, leaf area index and stem diameter (cm)). Yield and yield attributes were straw dry yield (ton/ha), full head dry weight (g), 1000 seeds weight (g), grain yield (ton/ha) and harvest Index (%). Data were analyzed using GEN STAT program. Means were separated using Least Significance Difference (LSD) at 5% probability level. Results showed that urea fertilizer significantly ($P \leq 0.05$) increased sorghum plant growth, yield attributes and grain yield compared to control in both seasons. Urea fertilizer significantly ($P \leq 0.05$) decreased days to 50% flowering and increased plant height (cm), straw dry yield (6.5 ton/ha) compared to control (4.7 ton/ha), 1000 seeds weight (g), grain yield (2.3 ton/ha) compared to control (1.9 ton/ha) and Harvest Index% for sorghum crop. Whereas; there was no significant effect amongst treatments on number of leaf per plant and stem diameter (cm). It was concluded that application of urea had a significant effect in most studied attributes in addition to shortening the maturity period. Therefore, to increase yield and yield components of Sorghum, namely cultivar

Remove the punctuation :[4GG]Comment

Which level of fertilizer :[5GG]Comment

Botana, in the study area during rainy season's application of urea is recommended.

Which treatment set-up did you recommend? Which is more general :[6GG]Comment

Key word: Urea fertilizer, Sorghum, Rainfed, North Kordofan State.

INTRODUCTION

Sorghum (*Sorghum bicolor* L. Moench) is the staple food crop of most people in the Sudan. In Sudan; sorghum is the most important cereal crop in terms of total acreage, production and consumption. Sorghum (Dura) is the main grain crop used by Sudanese in their daily diet (Mohammed *et al.*, 2020). It is mainly produced traditionally in rain fed areas and in modern agriculture in central clay plain Gezira.

General comment : [7GG]Comment
Why don't use paragraphs? .1
Lack of consistency?? .2
In my opinion, Botana, is the variety that you used in the current study, but you don't say .3
nothing about this variety, why?. So, you should put one paragraph which explains about this variety.
And why you used this variety? .4
The introduction is too short .5

Soil fertility is a major barrier affecting all aspects of crop production (Mbahand Mbagwu, 2006). ~~Area under irrigated sorghum is about 8% while 92% of it is under rain fed (Fadlelmula, 2009).~~ Low soil fertility is one of the main factors responsible for low productivity of crops in North Kordofan State. Soil fertility can be presumably enhanced by organic and inorganic fertilizers application (Khalid and Fadni, 2013). Soils in North Kordofan are predominantly sandy; these soils are characterized by low nutrient and water holding capacity, especially during hot summer. This often results in stunted plant growth, poor bloom or crop production. However, sandy soil also offers benefits, including excellent drainage, good air circulation and less pest incidence. By properly preparing and maintaining sandy soils, farmers enjoy these advantages while manipulating possible negative production factors.

Delete or move to the above : [8GG]Comment
unless it's not in its appropriate place

Soil fertility in smallholder farms is almost entirely dependent on locally available resources. Urea is a fertilizer material used for direct application to crops or in the preparation of blended fertilizers. Under most circumstances, it is equivalent to or superior than most other nitrogen sources. To understand the significance of urea, as opposed to other sources of nitrogen, one must examine the nature of

What does it mean? Do you mean: soil fertility improvement in the smallholder : [9GG]Comment

Observe these two sentences, and do you see any inconsistency?? : [10GG]Comment

commercial fertilizers and how they are blended. The current study is an attempt to evaluate the effect of urea nitrogen on growth and yield of Sorghum Cultivar (*Botana*) under rain-fed conditions in North Kordofan State Sheikan Locality.

MATERIALS AND METHODS

Study area

Sheikan locality lies in the center of the state of Kordofan. It covers an area of about 8312Km² (2 million acres), mostly useful for agriculture and grazing activities. Sheikan locality is located within the semi-desert zone where the average annual rainfall is between 250-450 mm/year during the rainy season (June to October).

Experimental Design and Treatments

~~For sorghum (cultivar *Botana*)~~, treatments assigned in a Randomized Complete Block Design (RCBD) with four replications for two consecutive seasons 2021/2022 and 2022/2023. The experiment was carried out at the premises of Faculty of Natural Resources and Environmental Studies, demonstration farm, in Sheikan locality, North Kordofan state. The plot area (2×5^m), the spacing for sorghum (50 × 50^{cm}) and the spacing between plot to plot (1^m). The first half dose was applied after three weeks from sowing, and the second one after a month from the first one. Seeds were sown at the depth of 2 cm at the middle of July for the two rainy seasons. The total rainfall for season 2021/2022 was (338.4mm/year) and was (317.9mm/year) for the second season 2022/2023. Rainfall amount and distribution were fairly similar in both seasons as shown in figure (1). The experiment consisted of three treatments Control (TR1), 40 kg urea/feddan (TR2) and 80 kg urea/feddan (TR3).

This study was conducted to :[11GG]Comment
evaluate the effect of urea nitrogen on growth and
yield components of Sorghum Cultivar (*Botana*)
under rain-fed conditions in North Kordofan State
Sheikan Locality

First and for most, "this :[12GG]Comment
study was carried out at the Sheikan locality. Then
you can say sth about that locality, unless and
otherwise, you couldn't jump directly to its
justification.

Based on the title, you :[13GG]Comment
should explain about the design and treatments
only.
So, this sub-heading is full of mistakes, and by
referring other literatures re-write it accordingly.

Move this to under study :[14GG]Comment
area

Move to other sub-heading. :[15GG]Comment
may be " site selection and preparation"

Move to under study area :[16GG]Comment




Figure 1: Rainfall amount (mm/year) during rainy seasons 2021/2022 and 2022/2023. Source: Elobeid Research Station.

Data Collection:

Plant growth parameters were days to 50% flowering, plant height (cm), number of leaf per plant, leaf area index and stem diameter (cm). Yield attributes were full head dry weight (g), straw dry yield (ton/ ha), grain yield (ton/ ha), 1000 – grain weight (g) and harvest index (%).

Data Analysis

Data was analyzed using GEN STAT program. Means were separated using Least Significance Difference (LSD) at 5% probability level. Based on the means, the percent change in each trait as compared to the control was determined.

Results

Sorghum Growth and Yield Parameters

Days to 50% flowering

Results showed that days to 50% flowering response decreased linearly (Table 1). All treatments showed significant ($P \leq 0.05$) early flowering effect in contrast to the

My buddy, you're sharing :[17GG]Comment knowledge to the scientific community. But, what's the rush?? Your title says "Result", but you discuss too. Take time to re-construct your findings!! Conversely, let's take it as "Result and discussion", which is not well discussed too.

control which displayed delayed flowering in both seasons. This observation suggests that the application of fertilizers decrease the number of days to the flowering stage implying earliness in maturity especially in such short rainy season. This could be attributed to the immediate accessibility of available nitrogen in the soil for plants, which plays a key role in cell division and enlargement (Muhammad *et al.* 2003).

Plant height (cm)

Table 1 illustrated that, the difference in values of plant height between urea and control was found to be significant at ($P \leq 0.05$) level at two seasons. The increase in height of plants obtained by the use of fertilizers might be due to the high stimulating effect of nitrogen on various physiological phases in cell division and cell elongation (Alim 2012). Urea Fertilizer provides adequate nitrogen levels during reproductive stage and support the formation of panicles and the filling of grains. This in turn was reflected in increasing plant growth parameters such as plant height and leaf area index. Fertilizer application in beans was also reported to improve soil conditions and supported better aeration to the plant roots and absorption of water and nutrients which was reflected in increasing plant growth parameters such as plant height (Manivannan *et al.*, 2009).

Number of leaf per plant

Effects of nitrogen application on the number of leaf per plant exhibited no significant differences amongst treatments in both seasons (Table 1). This could be due to that leaf number is dictated by the genetic setup of the plant (Meena and Mann. 2007). In other words, these results could be attributed to the length of the growth cycle of the cultivars which solely depends on the genetic makeup of the cultivar plants. However, increasing vegetative growth rate may lead to a reduction in nutrients in the leaf, by increasing their remobilization from leaf to grains, resulting in decreasing nutrients concentration in leaf with progress in plant growth (Al-Fadlly, 2011 and Mohammed, 2013).

Table 1: Effects of urea fertilizer on days to 50 % flowering (days) plant height (cm), number of leaf per plant, on sorghum plants for rainy seasons 2021/2022-2022/2023

Parameters						
Treatment	50% flowering		Plant height		No. of leaf /plant	
	Season 1	Season 2	Season 1	Season 2	Season 1	Season 2
TR1	49a	50a	65.1d	80.0b	10a	9a
TR2	44b	48a	69.4b	80.8b	9a	10a
TR3	44b	45b	60.8e	82.1a	9a	10a
Mean	45	48	65.1	80.9	9	10
LSD _(0.05)	1.4	3.4	7.9	7.3	1.1	0.6
SE \pm	0.43*	0.58*	1.52*	1.04ns	0.16ns	0.09ns
CV%	4.78	6.14	11.09	6.23	8.09	4.9

TR1: control

TR2: 40 kgurea / feddan

TR3: 80 kgurea/ feddan

Leaf area index (LAI)

Leaf area index was significantly ($P \leq 0.05$) increased due to urea fertilizer compared to control, in two seasons (table 2). These nutrients triggered the vigorous growth of plant leaves, thereby achieving higher LAI and further boosted the dry matter production and hastened the flowering and maturity period. Moreover, Aflakpule *et al.*, (2005), reported that N.P.K application increased growth and yield of sorghum. Similar results were stated by (Darwin *et al* 2018) who reported that the response of plants to the application of urea fertilizer gave a significant effect on leaf area index in sweet corn.

Stem diameter (cm)

The results of stem diameter showed that there were no significant differences ($P \leq 0.05$) between treatments at two seasons table (2). At stem diameter in sorghum

is thought to be genetically controlled. The nitrogen content also serves to spur the process of formation of the plant growth, because nitrogen is a nutrient-forming amino acid and proteins as raw material in the preparation of plant growth as explained by (Djunaed, 2009) in long beans. Stem diameter is dependent on the number of internodes for their length and this represents the genotypic characteristics of a variety (Abdul-Basit 2019).

Table 2: Effects of urea fertilizer on sorghum leaf area index and stem diameter for rainy seasons 2021/2022-2022/2023

Parameters				
Treatment	Leaf area index		Stem diameter	
	Season 1	Season 2	Season 1	Season 2
TR1	237.7c	226.4c	1.7a	1.6a
TR2	246.5b	259.0b	1.7a	1.6a
TR3	274.2a	287.3a	1.7a	1.7a
Mean	252.8	257.6	1.7	1.6
LSD	6.4	5.5	0.9	1.2
SE±	9.9**	7.9**	0.05ns	0.02ns
CV%	18.5	15.2	13.26	7.5

TR1: control

TR2: 40 kgurea / feddan

TR3: 80 kgurea/ feddan

Fullhead dry weight (g)

Table 3 showed that, applying urea fertilizer resulted in significantly ($P \leq 0.05$) higher full head dry weight as compared with control and the difference was, the highest weight was in season two (35.3 g) in contrast to control (21.4g) for the same season. The full head dry weight consistently increased at the two seasons. Increased in full head dry weight with an increased in nitrogen levels was obvious due to that fact, nitrogen promotes the plant growth manifesting positive effects on

full head dry weight. It was obvious that the increase in head dry weight is an essential factor for panicle weight and hence, increase in number and weight of grains per head and consequently on grain yield. An increase in the full dry head weight at higher nitrogen levels might be attributed to the lower competition for nutrients allowing the plants to accumulate more biomass with a higher capacity to convert more photosynthesis products into sink resulting in more grains (Kripa *et al* 2021).

Straw dry yield (ton/ ha)

Differences in straw dry weight between treatments were Significant ($P \leq 0.05$) in both seasons (Table 3). The highest weight of dry straw found when 80 kg urea/feddan applied in season one (5.6 ton/ha). The weight of dry straw can be used as a reference for measuring the growth of a plant as the fresh straw weight can rapidly change in a relatively short time due to loss of plant moisture content (Fitter and Hay 1994). Also, dry straw weight accumulates various food reserves, including protein, carbohydrates, and fats. According to (Moe *et al.* 2017) total dry matter accumulation in rice plants varied significantly under different levels of inorganic fertilizer. The effect of N fertilization on growth, yield and yield quality of two sorghum cultivars the high rate of N (80 IbN/feddan) significantly increased forage yield (fresh and dry) and crude fiber percent of *Abusabin* cultivar, (Zaid, 2004).

1000 seeds weight (g)

The result showed that application of 80 kg urea/feddan significantly ($P \leq 0.05$) gave the highest 1000 grain weight than other treatments in both seasons (table 4). These results are in agreement with findings of (ShuaibuYunusa *et al* 2018) who reported that the combination of organic and inorganic fertilizers increases the weight of 1000 seeds in sorghum. Other studies revealed significant increase in 1000-grain weight due to the addition of nitrogen alone (Mafongoya, 2006 and Chaudhary and Mehdi 2003). The effect of nitrogen fertilizer on 1000 grain yield of sorghum was reported to be significant only in the first sowing date (Azrag *et al* 2015).

Harvest index (%)

Results in Table 4 depicted that harvest index values were affected by application of 80kg urea/feddan which recorded the highest values of harvest index compared with other treatments at two seasons. Since there is a strong relationship between harvest index and nitrogen stock, the increase of harvest index in plants is due to increase of production and accumulation of assimilates during grain filling stage and assimilates remobilization after pollination. Previous results (Buah *et al.*, (2005) in pearl millet and the findings of (Muhammad Arif *et al.*, 2014) in rice, and Aflakpuet *et al.*, (2005), in millet and rice crops were consistent with the findings of this study.

Grain yield (ton/ha)

As presented in table 3, the effects of urea on grain yield of sorghum under study revealed that the application of 80 kg urea/feddan was found to produce significantly ($P \leq 0.05$) higher yield than other treatments used at the two seasons. Application of 80 kg urea/feddan increased the yield up to (2.6 ton/ha) compared to control (1.9 ton/ha) in season two. The higher grain yield obtained could be due to balance in the proportion of nutrients in adequate amount. This is supported by the finding of (Ibrahim and Hashim 2002) they further reported that application of urea to the soil significantly increased total nitrogen content of the top 30 cm of the soil. This is in conformity with the report of (Conley and Dunn, 2005 and Lafarge and Hamma, 2002) who indicated that the grain sorghum responded to inorganic fertilizer by increasing grain yield. It seems plausible that the increase in grain yield attributes unequivocally will be reflected in the final grain yield. The subtle differences between the two seasons could be attributed to rainfall amount and distribution added to differences in cultural practices.

In the current study :[18GG]Comment

????? :[19GG]Comment

Table 3: Effects of urea fertilizer on full head dry weight (g), straw dry yield (ton/ ha) and grain yield (ton/ ha) for rainy seasons 2021/2022- 2022/2023

Parameters						
Treatment	Head dry weight		Straw dry yield		Grain yield	
	Season 1	Season 2	Season 1	Season 2	Season 1	Season 2
TR1	17.7c	21.4c	4.7b	4.8b	1.9b	1.9b
TR2	25.2ab	32.1ab	5.1a	5.1a	2.1a	2.5a
TR3	26.0a	35.3a	5.6a	5.3a	2.3a	2.6a
Mean	23	29.6	5.1	5.1	2.1	2.3
LSD	0.6	3.9	1.9	1.7	0.5	0.2
SE±	0.27*	1.9*	0.52*	0.3*	0.08*	0.14*
CV%	23.13	19.2	18.67	22.9	17.07	16.1

TR1: control

TR2: 40 kgurea / feddan

TR3: 80 kgurea/ feddan

1000 seeds weight (g)

The result showed that application of 80 kg urea\ feddan was observed to have significantly ($P \leq 0.05$) the highest 1000 grain weight than other treatments in both seasons table 4. These results are in agreement with findings of (ShuaibuYunusa *et al* 2018) who reported that the combination of organic and inorganic fertilizers increases the weight of 1000 seeds in sorghum. Other studies revealed significant increase in 1000-grain weight due to the addition of nitrogen alone (Mafongoya, 2006 and Chaudhary and Mehdi 2003). The effect of nitrogen fertilizer on 1000 grain yield of sorghum was reported to be significant only in the first sowing date (Azrag *et al* 2015).

Harvest index (%)

Result in Table 4 show that harvest index values of sorghum plants were affected by different application rates of urea fertilizer. It was noticed that the harvest index

values were affected by 80kg urea/ feddan which recorded the highest values of harvest index compared with other treatments at two seasons. Since there is a strong relationship between harvest index and nitrogen stock, the increase of harvest index in plants is due to increase of production and synthesis of assimilates during grain filling stage and assimilates remobilization before pollination. The results of this study were also consistent with the findings of (Buah *et al.*, (2005) in pearl millet and the findings of (Muhammad Arif *et al.*,2014) in rice. (Aflakpul 2005), also reported that manure alone or mixed with N.P.K result in an increase in growth and yield of sorghum. Other studies have also shown the influence of different nutrient management practices on the harvest index of pearl millet (Bhanu, 2014).

Table 4:Effects of urea fertilizer on 1000 seeds weight (g) and harvest index (%) for rainy seasons 2021/2022- 2022/2023

Parameters				
Treatment	1000 seeds weight		Harvest index	
	Season 1	Season 2	Season 1	Season 2
TR1	10.5b	15.9b	30.2c	38.6c
TR2	13.2ab	18.5a	42.6b	41.9b
TR3	14.6a	18.7a	48.8a	44.0a
Mean	12.8	17.7	40.5	41.5
LSD	3.9	1.7	1.9	3.4
SE±	0.63*	0.32*	0.28**	1.9*
CV%	22.31	8.8	16.66	17.2

TR1: control

TR2: 40 kgurea / feddan

TR3: 80 kgurea/ feddan

CONCLUSIONS AND RECOMMENDATIONS

Application of 80 kg urea/feddan was found to significantly increase sorghum (*Botana* cultivar) plant height, leaf area index, stem diameter, full head dry weight (g), straw dry yield (ton/ ha), 1000 seeds weight (g), grain yield (ton/ ha), and harvest index. To increase productivity of Sorghum (cultivar *Botana*) in north kordofan State under rain-fed, urea fertilizer is recommended.

It's too short and I expected :[20GG]Comment that you will conclude your work according to your findings, but which is not

Which treatment do you :[21GG]Comment recommend?

References

1. Abdul-Basit Iddrisu Abdul-Rahman. 2019, Combined Effect of Organic and Inorganic Fertilizers on Growth of Rice Plants, Unu Land Restoration Training Programme, Final Project 2019.
2. Aflakpul GSK, Anchirinah VM and Asumadu H. 2005, Respond of quality protein maize hybrid to N supply and plant density in the forest zone of Ghana, Tropical. Science, 45, 3-7.
3. Al Fadlly, J.T.M. 2011, Effect of organic and mineral fertilization on growth and yield of potato crop. M. Sc. Thesis. Coll. Agric. Univ. Baghdad, Iraq. (in Arabic).
4. Alim M. 2012, Effect of organic and inorganic sources and doses of nitrogen fertilizer on the yield of boro rice. Journal of Environmental Science and Natural Resources 5:273–282.
5. Azrag, A.A.D, Dagash, Y.M. I and Yagoub, S.O. 2015, Effect of Sowing Date and Nitrogen Fertilizer Rate on Yield of Sorghum (*Sorghum bicolor* L.) and Nitrogen Use Efficiency. Journal of Agricultural and veterinary Sciences. Vol 16 No (1).
6. Bhanu Prasad Reddy S. 2014, Effect of nitrogen and potassium on yield and quality of pearl millet, thesis submitted to the acharya n.g. ranga agricultural university in partial fulfillment of the requirements for the award of the degree of master of science in agriculture (soil science and agricultural

Use published only?? :[22GG]Comment

chemistry) chairperson: dr. K.v. Naga madhuri department of soil science and agricultural chemistry.

7. Buah S and Maranville G. 2005), Response of nitrogen use efficient sorghum to nitrogen fertilizer. *Journal of Plant Nutrition*, 28:2303-2318.
8. Chaudhary EH, Ranjha AM, Gill AM and Mehdi SM. 2003, phosphorus requirement of maize in relation to soil characteristic. *International journal of agricultura biology*, 5(4), 625-629.
9. Conley SP, Steven WG and Dunn DD. 2005, Grain sorghum responds to row spacing, plant density and planter skips. *Crop management*, 5, 56- 59.
10. Darwin Habinsarand Pangaribuan, Kus Hendarto, Sheilla R. Elzhivago, Ade Yulistiani. 2018. The Effect of Organic Fertilizer and Urea Fertilizer on Growth, Yield and Quality of Sweet Corn and Soil Health. *Asian J Agri & Biol*. 2018;6(3):335-344.
11. Djunaed, A. 2009, Effect of Botashi Fertilizer Dose on Growth and Production Long Bean (*Vigna Sinensis* L). *Agrovigor*, 2:42-46.
12. Fadlelmula, A. A, 2009. Effect of sowing date on the incidence of sorghum midge, *Stenodiplosis sorghicola* Coq.), (Cecidomyiidae Diptera): in sorghum (*Sorghum bicolor* L. Moench) at Damazin rain-fed area, Sudan
13. Fitter A H and Hay R K M. 1994, *Plant Environmental Physiology* Translator: Andani S and Purbayanti E D (Yogyakarta: Gadjah Mada University Press) of Agriculture and Forestry, 2(2), 56-60.
14. Ibrahim, S.H.; Moamed, A.B.; Osman, H.N., And Hashim, A.A. 2002, Fertilizing with Urea and FYM For Higher Cotton Yields and Better Soil Conditions in The Gezira. Paper Presented in The National Crop Husbandry Committee Meeting, 2002, Agricultural Research Corporation (ARC), Wad Medani Sudan.
15. Khalid Ibrahimh. M. And Fadni O.A.S. 2013, Effect of Organic Fertilizers Application on Growth, Yield and Quality of Tomatoes in North Kordofan

Too old replace with latest :[23GG]Comment one!

(Sandy Soil) Western Sudan. Greener Journal of Agricultural Sciences ISSN: 2276-7770; ICV: 6.15. Vol. 3(4), 299-304, April 2013.

16. Kripa Adhikari¹, Sudip Bhandari, Krishna Aryal, Mohan Mahato And Jiban Shrestha. 2021, Effect of Different Levels of Nitrogen on Growth And Yield Of Hybrid Maize (*Zea Mays* L.) Varieties, Journal of Agriculture And Natural Resources (2021) 4(2): 48-62 ISSN: 2661-6270 (Print), ISSN: 2661-6289 (Online) DOI: <https://doi.org/10.3126/Janr.V4i2.33656>.
17. Lafarge TA And Hamma GA. 2002, Tillering in Grain Sorghum Over A Wide Range of Population Densities. Modeling Dynamic of Tiller Density. Bot., Journal of Agronomy, 6(9), 99-110.
18. Mafongoya PL, Bationo A, Kihara J and Swaswa B. 2006, Appropriate Technologies to Replenish Soil Fertility in Southern Africa. Nutrition Cycle Agroecology, 76, 137 - 151.
19. Manivannan, S.; M. Balamurugan; K. Parthasarathi; G. Gunasekharan And R. Ranganathan. 2009, Effect of Vermicompost on Soil Fertility and Crop Productivity Beans (*Phaseolus Vulgaris*) J. Environ, 30: 275-281.
20. Mbah CN, Mbagwu JS, 2006. Effect of Animal Waste on Physicochemical Properties of a Dystric Leptosol And Maize Yield in Southeastern Nigeria. Nig J Soil Sci; 1: 290 -305.
21. Meena L, Mann J. 2007, Response of Forage Sorghum (*Sorghum Bicolor*) Varieties to Different Levels of Nitrogen with and Without Farmyard Manure in A Semiarid Region. International Journal of Tropical Agriculture, 25: 105-109.
22. Moe K, Mg KW, Win KK, Yamakawa T. 2017, Combined Effect of Organic Manures and Inorganic Fertilizers on The Growth and Yield of Hybrid Rice (Palethwe-1). American Journal of Plant Sciences 8:1022–1042
23. Mohammed Abdalla Elsheikh¹, Mubarak Abdelrahman Abdalla¹ And Abdel Moneim Mohammed Ahmed El Tilib 2020, Advances in Soil Fertility Research in Sudan, Desertification and Desert Cultivation Studies Institute,

- University of Khartoum, Sudan, Sudan J. Des. Res. Vol. 12 (1):96-129, 2020, At: <https://www.researchgate.net/publication/343862739>.
24. Mohammed, E.J. 2013, Evaluation the Effect of Organic and Mineral Fertilization on Some Soil Properties, Growth and Yield of Snake Cucumber (*Cucumis Melo* Var. *Flexuosus* nauds). M. Sc. Thesis. Coll. Agric. Univ. Basrah, Iraq. (In Arabic).
25. Muhammad Arif, Muhammad Tasneem, Fiaz Bashir, Ghulam Yaseen, And Rana Muhammad Iqbal. 2014, Effect of Integrated Use of Organic Manures and Inorganic Fertilizers on Yield and Yield Components of Rice. Chakkanwali Reclamation Research Station, Directorate of Land Reclamation, Punjab, Pakistan. J. Agric. Res., 2014, 52(2)
26. Muhammad U, Ehsan U, Ejaz AW, Muhammad FAL. 2003, Effect of Organic and Inorganic Manures on Growth and Yield of Rice Variety "Basmati – 2000". International Journal of Agriculture & Biology 4:481–483
27. Shuaibu Yunusa Muhammad, Bala Rashida Abdulmumini, Kawure Sani And Shuaibu Zaharaddeen. 2018, Effect of Organic and Inorganic Fertilizer on The Growth and Yield of Sorghum (*Sorghum Bicolor* (L.) Moench) In Bauchi State, Nigeria. Available Online At GSC, Online Press Directory GSC Biological and Pharmaceutical Sciences, E-ISSN: 2581-3250, CODEN (USA): GBPSC2, Journal Homepage: <https://www.gsc-onlinepress.com/journals/gscbps>.
28. Siavoshi M, Nasiri A, Lawere SL .2011, Effect of Organic Fertilizer on Growth and Yield Components in Rice (*Oryza Sativa* L.). Journal of Agricultural Science 3:217-224.
29. Zaid, G.A.A. 2004. Effect of Nitrogen and Sulfur Fertilization on Growth, Yield, Quality, Nitrogen and Prussic Acid Content of Tow Forage Sorghum Cultivars. M.Sc. (Agric) Thesis, University of Khartoum.

UNDER PEER REVIEW