

## INDICES OF LINE GRAPH OF $\mathcal{R}(\mathcal{G})$

**ABSTRACT.** The field of graph theory facilitates in the study of chemical, physical, and biological characteristics of chemical substances. The properties and reactivity of chemical compounds can be predicted by examining the graph of chemical structure and its topological indices. Topological index of graph is a powerful tool for sidestepping costly and tedious experiments which should be done in Laboratory. More than 140 topological indices have been defined so far and more studies are still going on in this area. In this study, we compute a few topological indices of line graphs of triangular graphs of specific families of graphs.

Let  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$  be a graph with node set  $\mathcal{V}$  and arc set  $\mathcal{E}$ . The number of nodes in  $\mathcal{G}$  is denoted by  $|\mathcal{V}| = \eta$  and the number of arcs in  $\mathcal{G}$  is denoted by  $|\mathcal{E}| = \zeta$ . The degree of the node  $x \in \mathcal{V}$  is denoted by  $\kappa_x$ , is the number of arcs incident on  $x$ . The arc connecting the nodes  $x$  and  $y$  is denoted by  $e = xy$ , where  $e \in \mathcal{E}$ .

The Triangle graph  $\mathcal{R}(\mathcal{G})$  is obtained from  $\mathcal{G}$  by adding one node corresponding to each arc and is joined to the end nodes of the corresponding arc. Line graph  $\mathcal{L}(\mathcal{G})$  is obtained from  $\mathcal{G}$  by taking the arcs of  $\mathcal{G}$  as the nodes and is joined by an arc in  $\mathcal{L}(\mathcal{G})$  if they have common node in  $\mathcal{G}$ .

The numerical value that results from a mathematical analysis of a graph is called a topological index. The properties and activities of chemical substances can be predicted with the use of this mathematical research. Finding the topological indices is a crucial field of study. There are three categories for topological indices: degree-based, distance-based, combination of degree-and distance-based, and counting-related topological indices. Degree-based topological indices are among these indicators that are crucial for predicting the characteristics and actions of chemical substances. Known topological indices include the oldest, Zagreb indices, the most recent Gourava indices, Randić' index, Atom bond index, Harmonic index, Sum connectivity index, and Sombor index.

In this paper, we calculated certain topological indices of line graph of triangle graph of a regular graph, Path  $\mathcal{P}_\eta$ , Cycle  $\mathcal{C}_\eta$ , Complete  $\mathcal{K}_\eta$  and Ladder graph  $\mathcal{L}_\eta$

Name	Topological Indices
Atom bond Connectivity Index [2]	$ABC(\mathcal{G}) = \sum_{xy \in \mathcal{E}} \sqrt{\frac{\kappa_x + \kappa_y - 2}{\kappa_x \kappa_y}}$
Randic Index[3]	$\mathcal{RA}(\mathcal{G}) = \sum_{xy \in \mathcal{E}} \sqrt{\frac{1}{\kappa_x \kappa_y}}$
sum-Connectivity Index [4]	$\chi(\mathcal{G}) = \sum_{xy \in \mathcal{E}} \sqrt{\frac{1}{\kappa_x + \kappa_y}}$
Geometric Arithmetic Index[5]	$\mathcal{GA}(\mathcal{G}) = \sum_{xy \in \mathcal{E}} \frac{2\sqrt{\kappa_x \kappa_y}}{\kappa_x + \kappa_y}$
Sombor Index [1]	$\mathcal{SO}(\mathcal{G}) = \sum_{xy \in \mathcal{E}} \sqrt{\kappa_x^2 + \kappa_y^2}$
First Zagreb Index [6]	$\mathcal{M}_1(\mathcal{G}) = \sum_{xy \in \mathcal{E}} \kappa_x + \kappa_y$
Second Zagreb Index [6]	$\mathcal{M}_2(\mathcal{G}) = \sum_{xy \in \mathcal{E}} \kappa_x \kappa_y$
First Gourava index [7]	$\mathcal{GO}_1(\mathcal{G}) = \sum_{xy \in \mathcal{E}} [(\kappa_x + \kappa_y) + \kappa_x \kappa_y]$
second Gourava index [7]	$\mathcal{GO}_2(\mathcal{G}) = \sum_{xy \in \mathcal{E}} [(\kappa_x + \kappa_y) \kappa_x \kappa_y]$

### 1. Topological indices of Line graph of Triangle graph of a regular graph

**Theorem 1.1.** *Let  $\mathcal{G}$  be a  $\kappa$  - regular graph with vertex set  $\mathcal{V}$  and edge set  $\mathcal{E}$  containing  $\eta$  and  $\zeta$  elements respectively. Then the topological indices of  $\mathcal{L}(\mathcal{R}(\mathcal{G}))$  is given by*

$$\begin{aligned}
(1) \quad ABC(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \kappa\zeta \sqrt{\frac{2(3\kappa-2)}{\kappa(2\kappa-1)}} + \frac{\zeta \sqrt{2(2\kappa-1)}}{2} + \frac{(2\kappa^2\eta - \zeta(3\kappa+1)) \sqrt{2}\sqrt{4\kappa-3}}{2(2\kappa-1)} \\
(2) \quad \mathcal{RA}(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \frac{\kappa\zeta}{\sqrt{\kappa(2\kappa-1)}} + \frac{\zeta}{2} + \frac{2\kappa^2\eta - \zeta(3\kappa+1)}{2(2\kappa-1)} \\
(3) \quad \chi(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \frac{\sqrt{2}\kappa\zeta}{\sqrt{3\kappa-1}} + \frac{\zeta\sqrt{\kappa}}{2} + \frac{2\kappa^2\eta - \zeta(3\kappa+1)}{2\sqrt{2\kappa-1}} \\
(4) \quad \mathcal{GA}(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \frac{4\kappa\zeta\sqrt{\kappa(2\kappa-1)}}{3\kappa-1} + 2\kappa^2\eta - 2\kappa\zeta - \zeta \\
(5) \quad \mathcal{SO}(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= 4\kappa\zeta\sqrt{5\kappa^2-4\kappa+1} + 2\sqrt{2}(\kappa^2\zeta + (2\kappa-1)(2\kappa^2\eta - \zeta(3\kappa+1))) \\
(6) \quad \mathcal{M}_1(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= 4\kappa\zeta(4\kappa-1) + 4(2\kappa-1)(2\kappa^2\eta - \zeta(3\kappa+1)) \\
(7) \quad \mathcal{M}_2(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= 8\kappa^2\zeta(2\kappa-1) + 4\kappa^3\zeta + 4(2\kappa^2\eta - \zeta(3\kappa+1))(2\kappa-1)^2 \\
(8) \quad \mathcal{GO}_1(\mathcal{G}) &= 4\kappa[4\kappa(\kappa\eta - \zeta)(2\kappa-1) + \zeta(\kappa^2+1)] \\
(9) \quad \mathcal{GO}_2(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= 16\kappa^2\zeta(2\kappa-1)(3\kappa-1) + 16\kappa^4\zeta + 8(2\kappa-1)[2\kappa^2\eta - \zeta(3\kappa+1)]
\end{aligned}$$

*Proof.* Let  $\mathcal{G}$  be a  $\kappa$  - regular graph with  $\eta$  vertices and  $\zeta$  edges. Then the  $\mathcal{R}(\mathcal{G})$  has  $\eta + \zeta$  vertices and  $3\zeta$  edges with  $\eta$  vertices of degree  $2\kappa$  and  $\zeta$  vertices of degree 2. So the graph  $\mathcal{L}(\mathcal{R}(\mathcal{G}))$  has  $3\zeta$  vertices and  $2\kappa^2\eta - \zeta$  edges with  $2\zeta$  vertices of degree  $2\kappa$  and  $\zeta$  vertices of degree  $4\kappa - 2$ .

**The edge partition of  $\mathcal{L}(\mathcal{R}(\mathcal{G}))$** 

Degree of end vertices	No. of edges
$(2\kappa, 2\kappa)$	$\kappa\zeta$
$(4\kappa - 2, 2\kappa)$	$2\kappa\zeta$
$(4\kappa - 2, 4\kappa - 2)$	$2\kappa^2\eta - \zeta(3\kappa + 1)$

$$\begin{aligned}
ABC(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} \sqrt{\frac{\kappa_x + \kappa_y - 2}{\kappa_x \kappa_y}} \\
&= 2\kappa\zeta \sqrt{\frac{4\kappa - 2 + 2\kappa - 2}{(2\kappa)(4\kappa - 2)}} + \kappa\zeta \sqrt{\frac{2\kappa + 2\kappa - 2}{(2\kappa)(2\kappa)}} + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa + 1)) \sqrt{\frac{4\kappa - 2 + 4\kappa - 2 - 2}{(4\kappa - 2)(4\kappa - 2)}} \\
&= \kappa\zeta \sqrt{\frac{2(3\kappa - 2)}{\kappa(2\kappa - 1)}} + \frac{\zeta\sqrt{2(2\kappa - 1)}}{2} + \frac{(2\kappa^2\eta - \zeta(3\kappa + 1))\sqrt{2}\sqrt{4\kappa - 3}}{2(2\kappa - 1)}
\end{aligned}$$

$$\begin{aligned}
\mathcal{RA}(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} \sqrt{\frac{1}{\kappa_x \kappa_y}} \\
&= 2\kappa\zeta \sqrt{\frac{1}{(2\kappa)(4\kappa - 2)}} + \kappa\zeta \sqrt{\frac{1}{(2\kappa)(2\kappa)}} + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa + 1)) \sqrt{\frac{1}{(4\kappa - 2)(4\kappa - 2)}} \\
&= \frac{\kappa\zeta}{\sqrt{\kappa(2\kappa - 1)}} + \frac{\zeta}{2} + \frac{2\kappa^2\eta - \zeta(3\kappa + 1)}{2(2\kappa - 1)}
\end{aligned}$$

$$\begin{aligned}
\chi(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} \sqrt{\frac{1}{\kappa_x + \kappa_y}} \\
&= 2\kappa\zeta \sqrt{\frac{1}{(2\kappa) + (4\kappa - 2)}} + \kappa\zeta \sqrt{\frac{1}{(2\kappa) + (2\kappa)}} + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa + 1)) \sqrt{\frac{1}{(4\kappa - 2) + (4\kappa - 2)}} \\
&= \frac{\sqrt{2}\kappa\zeta}{\sqrt{3\kappa - 1}} + \frac{\zeta\sqrt{\kappa}}{2} + \frac{2\kappa^2\eta - \zeta(3\kappa + 1)}{2\sqrt{2\kappa - 1}}
\end{aligned}$$




$$\begin{aligned}
\mathcal{GA}(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} 2 \frac{\sqrt{\kappa_x \kappa_y}}{\kappa_x + \kappa_y} \\
&= (2\kappa\zeta) 2 \frac{\sqrt{2\kappa(4\kappa-2)}}{(2\kappa) + (4\kappa-2)} + 2\kappa\zeta \frac{\sqrt{(2\kappa)(2\kappa)}}{(2\kappa) + (2\kappa)} + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa+1)) 2 \sqrt{\frac{(4\kappa-2)(4\kappa-2)}{(4\kappa-2) + (4\kappa-2)}} \\
&= \frac{4\kappa\zeta \sqrt{\kappa(2\kappa-1)}}{3\kappa-1} + 2\kappa^2\eta - 2\kappa\zeta - \zeta \\
\mathcal{SO}(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} \sqrt{\kappa_x^2 + \kappa_y^2} \\
&= 2\kappa\zeta \sqrt{(2\kappa)^2 + (4\kappa-2)^2} + \kappa\zeta \sqrt{(2\kappa)^2 + (2\kappa)^2} \\
&\quad + (2\kappa^2\eta - \zeta(3\kappa+1)) \sqrt{(4\kappa-2)^2 + (4\kappa-2)^2} \\
&= 4\kappa\zeta \sqrt{5\kappa^2 - 4\kappa + 1} + 2\sqrt{2} (\kappa^2\zeta + (2\kappa-1)(2\kappa^2\eta - \zeta(3\kappa+1))) \\
\mathcal{M}_1(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} \kappa_x + \kappa_y \\
&= 2\kappa\zeta (4\kappa-2 + 2\kappa) + \kappa\zeta (2\kappa + 2\kappa) + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa+1)) (4\kappa-2 + 4\kappa-2) \\
&= 4\kappa\zeta (4\kappa-1) + 4(2\kappa-1) (2\kappa^2\eta - \zeta(3\kappa+1)) \\
\mathcal{M}_2(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} \kappa_x \kappa_y \\
&= 2\kappa\zeta [(4\kappa-2)(2\kappa)] + \kappa\zeta [(2\kappa)(2\kappa)] + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa+1)) [(4\kappa-2)(4\kappa-2)] \\
&= 8\kappa^2\zeta (2\kappa-1) + 4\kappa^3\zeta + 4(2\kappa^2\eta - \zeta(3\kappa+1)) (2\kappa-1)^2 \\
\mathcal{GO}_1(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} [(\kappa_x + \kappa_y) + \kappa_x \kappa_y] \\
&= 2\kappa\zeta [(4\kappa-2) + 2\kappa + 2\kappa(4\kappa-2)] + \kappa\zeta [(2\kappa) + (2\kappa) + (2\kappa)^2] + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa+1)) [(4\kappa-2) + (4\kappa-2) + (4\kappa-2)^2] \\
&= 4\kappa [4\kappa(\kappa\eta - \zeta)(2\kappa-1) + \zeta(\kappa^2+1)] \\
\mathcal{GO}_2(\mathcal{L}(\mathcal{R}(\mathcal{G}))) &= \sum_{xy \in \mathcal{E}} \kappa_x \kappa_y (\kappa_x + \kappa_y) \\
&= 2\kappa\zeta [(4\kappa-2)^2(2\kappa) + (4\kappa-2)(2\kappa)^2] + \kappa\zeta [(2\kappa)(2\kappa)^2 + (2\kappa)^2(2\kappa)] + \\
&\quad (2\kappa^2\eta - \zeta(3\kappa+1)) [(4\kappa-2)(4\kappa-2)^2 + (4\kappa-2)^2(4\kappa-2)] \\
&= 16\kappa^2\zeta (2\kappa-1)(3\kappa-1) + 16\kappa^4\zeta + 8(2\kappa-1) [2\kappa^2\eta - \zeta(3\kappa+1)]
\end{aligned}$$

□

2. Topological indices of  $\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))$ 

**Theorem 2.1.** Let  $\mathcal{P}_\eta$  be a path with  $\eta$  vertices. Then the topological indices of  $\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))$  is given by

- (1)  $ABC(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = \frac{1}{6} [12\sqrt{2} + 3\sqrt{6}(2\eta - 1) + 4\sqrt{3}(2\eta - 5) + \sqrt{10}(\eta - 4)]$
- (2)  $\mathcal{RA}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = \frac{1}{12} [3\sqrt{2} + 3(2\eta - 1) + 2\sqrt{6}(2\eta - 5) + 2(\eta - 4)]$
- (3)  $\chi(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = \frac{1}{30} [20\sqrt{6} + 30\sqrt{2}(2\eta - 1) + 6\sqrt{10}(2\eta - 5) + 5\sqrt{3}(\eta - 4)]$
- (4)  $\mathcal{GA}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = \frac{1}{15} [80\sqrt{3} + 30\sqrt{2}(2\eta - 1) + 24\sqrt{15}(2\eta - 5) + 30\sqrt{3}(\eta - 4)]$
- (5)  $\mathcal{SO}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = \eta(14\sqrt{2} + 8\sqrt{13}) + 8\sqrt{5} - 28\sqrt{2} - 20\sqrt{13}$
- (6)  $\mathcal{M}_1(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = 4(17\eta - 33)$
- (7)  $\mathcal{M}_2(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = 4(41\eta - 92)$
- (8)  $\mathcal{GO}_1(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = 4(58\eta - 125)$
- (9)  $\mathcal{GO}_2(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = 16(73\eta - 254)$

FIGURE 1.  $\mathcal{P}_3$ FIGURE 2.  $\mathcal{R}(\mathcal{P}_3)$ FIGURE 3.  $\mathcal{L}(\mathcal{R}(\mathcal{P}_3))$ 

*Proof.* Let  $\mathcal{P}_\eta$  be a path with  $\eta$  vertices. Then  $\mathcal{R}(\mathcal{P}_\eta)$  has  $2\eta - 1$  vertices and  $3(\eta - 1)$  edges. Therefore  $\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))$  has  $3(\eta - 1)$  vertices and  $7\eta - 11$  edges.

**The edge partition of  $\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))$** 

Degree of end vertices	No. of edges
(2, 2)	4
(4, 4)	$2\eta - 1$
(4, 6)	$4\eta - 10$
(6, 6)	$\eta - 4$



$$\begin{aligned}
\mathcal{ABC}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4\sqrt{\frac{2+4-2}{(2)(4)}} + (2\eta-1)\sqrt{\frac{4+4-2}{(4)(4)}} + (4\eta-10)\sqrt{\frac{4+6-2}{(4)(6)}} \\
&\quad + (\eta-4)\sqrt{\frac{6+6-2}{(6)(6)}} \\
&= \frac{1}{6} \left[ 12\sqrt{2} + 3\sqrt{6}(2\eta-1) + 4\sqrt{3}(2\eta-5) + \sqrt{10}(\eta-4) \right] \\
\mathcal{RA}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4\sqrt{\frac{1}{(2)(4)}} + (2\eta-1)\sqrt{\frac{1}{(4)(4)}} + (4\eta-10)\sqrt{\frac{1}{(4)(6)}} + (\eta-4)\sqrt{\frac{1}{(6)(6)}} \\
&= \frac{1}{12} \left[ 3\sqrt{2} + 3(2\eta-1) + 2\sqrt{6}(2\eta-5) + 2(\eta-4) \right] \\
\chi(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4\sqrt{\frac{1}{2+4}} + (2\eta-1)\sqrt{\frac{1}{4+4}} + (4\eta-10)\sqrt{\frac{1}{4+6}} + (\eta-4)\sqrt{\frac{1}{6+6}} \\
&= \frac{1}{30} \left[ 20\sqrt{6} + 30\sqrt{2}(2\eta-1) + 6\sqrt{10}(2\eta-5) + 5\sqrt{3}(\eta-4) \right] \\
\mathcal{GA}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4(2)\sqrt{\frac{(2)(4)}{2+4}} + (2\eta-1)(2)\sqrt{\frac{(4)(4)}{4+4}} + (4\eta-10)(2)\sqrt{\frac{(4)(6)}{4+6}} \\
&\quad + (\eta-4)(2)\sqrt{\frac{(6)(6)}{6+6}} \\
&= \frac{1}{15} \left[ 80\sqrt{3} + 30\sqrt{2}(2\eta-1) + 24\sqrt{15}(2\eta-5) + 30\sqrt{3}(\eta-4) \right] \\
\mathcal{SO}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4\sqrt{2^2+4^2} + (2\eta-1)\sqrt{4^2+4^2} + (4\eta-10)\sqrt{4^2+6^2} + (\eta-4)\sqrt{6^2+6^2} \\
&= \eta \left( 14\sqrt{2} + 8\sqrt{13} \right) + 8\sqrt{5} - 28\sqrt{2} - 20\sqrt{13} \\
\mathcal{M}_1(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4(2+4) + (2\eta-1)(4+4) + (4\eta-10)(4+6) + (\eta-4)(6+6) \\
&= 4(17\eta-33) \\
\mathcal{M}_2(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4(2.4) + (2\eta-1)(4.4) + (4\eta-10)(4.6) + (\eta-4)(6.6) \\
&= 4(41\eta-92) \\
\mathcal{GO}_1(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4(2+4+2.4) + (2\eta-1)(4+4+4.4) + (4\eta-10)(4+6+4.6) \\
&\quad + (\eta-4)(6+6+6.6) \\
&= 4(58\eta-125) \\
\mathcal{GO}_2(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) &= 4(2.4)(2+4) + (2\eta-1)(4.4)(4+4) + (4\eta-10)(4.6)(4+6) \\
&\quad + (\eta-4)(6.6)(6+6) \\
&= 16(73\eta-254)
\end{aligned}$$

□

### 3. Topological indices of $\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))$

**Theorem 3.1.** *Let  $\mathcal{C}_\eta$  be a cycle with  $\eta$  vertices. Then the topological indices of  $\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))$  is given by*

- (1)  $ABC(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = \frac{\eta}{6} (3\sqrt{6} + 8\sqrt{3} + \sqrt{10})$
- (2)  $\mathcal{RA}(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = \frac{\eta}{3} (2 + \sqrt{6})$
- (3)  $\chi(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = \frac{\eta}{30} (30\sqrt{2} + 12\sqrt{10} + 5\sqrt{3})$
- (4)  $\mathcal{GA}(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = \eta (4\sqrt{2} + 16\sqrt{15} + 2\sqrt{3})$
- (5)  $\mathcal{SO}(\mathcal{L}(\mathcal{R}(\mathcal{P}_\eta))) = \eta (14\sqrt{2} + 8\sqrt{13})$
- (6)  $\mathcal{M}_1(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = 68\eta$
- (7)  $\mathcal{M}_2(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = 164\eta$
- (8)  $\mathcal{GO}_1(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = 232\eta$
- (9)  $\mathcal{GO}_2(\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))) = 1648\eta$

FIGURE 4.  $\mathcal{R}(\mathcal{C}_4)$ FIGURE 5.  $\mathcal{L}(\mathcal{R}(\mathcal{C}_4))$

*Proof.* Let  $\mathcal{C}_\eta$  be a cycle with  $\eta$  vertices. Then  $\mathcal{R}(\mathcal{C}_\eta)$  has  $2\eta$  vertices and  $3\eta$  edges. Therefore  $\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))$  has  $3\eta$  vertices and  $7\eta$  edges.

***The edge partition of  $\mathcal{L}(\mathcal{R}(\mathcal{C}_\eta))$***

Degree of end vertices	No. of edges
(4, 4)	$2\eta$
(4, 4)	$4\eta$
(6, 6)	$\eta$

The proof is similar as above. □

#### 4. Topological indices of $\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))$

**Theorem 4.1.** *Let  $\mathcal{K}_\eta$  be a complete with  $\eta$  vertices . Then the topological indices of  $\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))$  is given by*

- (1)  $ABC(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = \eta C_2 \left[ \frac{\sqrt{4\eta-6}}{2} + \sqrt{\frac{2(3\eta-5)(\eta-1)}{(2\eta-3)}} + \frac{\eta-2}{(2\eta-3)} \sqrt{2(4\eta-7)} \right]$
- (2)  $RA(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = \eta C_2 \left[ \frac{1}{2} + \sqrt{\frac{\eta-1}{2\eta-3}} + \frac{\eta-2}{2(\eta-3)} \right]$
- (3)  $\chi(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = \eta C_2 \left[ \frac{\sqrt{\eta-1}}{2} + \frac{\sqrt{2}(\eta-1)}{\sqrt{(3\eta-4)}} + \frac{\eta-2}{\sqrt{(2\eta-3)}} \right]$
- (4)  $GA(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = 2(\eta C_2) \left[ \frac{2\eta-3}{2} + 2\frac{\eta-1}{3\eta-4} \sqrt{(\eta-1)(\eta-3)} \right]$
- (5)  $SO(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = (2\sqrt{2})\eta C_2 \left[ (3\eta^2 - 9\eta + 7) + \sqrt{2}(\eta-1) \sqrt{5\eta^2 - 14\eta + 10} \right]$
- (6)  $M_1(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = 4(\eta C_2) (6\eta^2 - 16\eta + 11)$
- (7)  $M_2(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = 4(\eta C_2) (9\eta^3 - 37\eta^2 + 52\eta - 25)$
- (8)  $GO_1(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = 4(\eta-1)(\eta C_2) (9\eta^2 - 22\eta + 14)$
- (9)  $GO_2(\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))) = 16(\eta C_2) (15\eta^4 - 85\eta^3 + 184\eta^2 - 168\eta + 67)$

*Proof.* Let  $\mathcal{K}_\eta$  be a complete graph with  $\eta$  vertices. Then  $\mathcal{R}(\mathcal{K}_\eta)$  has  $\eta + \eta C_2$  vertices and  $3(\eta C_2)$  edges. Therefore  $\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))$  has  $3(\eta C_2)$  vertices and  $\eta C_2(4\eta - 5)$  edges

***The edge partition of  $\mathcal{L}(\mathcal{R}(\mathcal{K}_\eta))$***

Degree of end vertices	No. of edges
$(2\eta-2, 2\eta-2)$	$(\eta-1)\eta C_2$
$(2\eta-2, 4\eta-6)$	$2(\eta-1)\eta C_2$
$(4\eta-6, 4\eta-6)$	$(\eta-2)\eta C_2$



The proof is similar as above.

□

### 5. Topological indices of $\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))$



**Theorem 5.1.** *Let  $\mathcal{L}_\eta$  be a Ladder graph. Then the topological indices of  $\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))$  is given by*

$$\begin{aligned}
 (1) \quad ABC(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= \eta \left[ \frac{3\sqrt{10}}{2} + \frac{3\sqrt{210}}{5} + \frac{9\sqrt{2}}{5} \right] + 8 + 3\sqrt{6} + \\
 &\quad 4\sqrt{3} + \frac{18\sqrt{5}}{5} - \frac{10\sqrt{10}}{3} - \frac{8\sqrt{210}}{5} - 6\sqrt{2}. \\
 (2) \quad \mathcal{RA}(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= \frac{\eta}{10} \left[ 6\sqrt{15} + 21 \right] + \sqrt{2} + \frac{4}{3}\sqrt{3} + \frac{2}{5}\sqrt{5} + \sqrt{6} - \frac{8}{5}\sqrt{15} - \frac{23}{6}. \\
 (3) \quad \chi(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= \frac{\sqrt{5}\eta}{10} \left[ 6 + 9\sqrt{5} + 3\sqrt{15} \right] + \frac{17}{6}\sqrt{2} + \frac{6}{5}\sqrt{10} + \frac{8}{7}\sqrt{14} - 2\sqrt{3} - 2\sqrt{5} - 12. \\
 (4) \quad \mathcal{GA}(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= \frac{3\eta}{2} \left[ 10 + 3\sqrt{5} \right] + \frac{16}{3}\sqrt{2} + \frac{64}{7}\sqrt{3} + \frac{32}{9}\sqrt{5} \\
 &\quad + \frac{24}{5}\sqrt{6} - 12\sqrt{15} - 37. \\
 (5) \quad \mathcal{SO}(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= 6\eta \left[ 19\sqrt{2} + 6\sqrt{34} \right] + 8 \left[ 4\sqrt{5} + 3\sqrt{13} + 2\sqrt{41} - 12\sqrt{34} - 56\sqrt{2} \right] \\
 (6) \quad \mathcal{M}_1(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= 4\eta (129\eta - 204) \\
 (7) \quad \mathcal{M}_2(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= 4\eta (501\eta - 888) \\
 (8) \quad \mathcal{GO}_1(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= 8 (315\eta - 313) \\
 (9) \quad \mathcal{GO}_2(\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))) &= 16 (2073\eta - 4108)
 \end{aligned}$$

*Proof.* The ladder graph  $\mathcal{L}_\eta$  has  $2\eta$  vertices and  $3\eta - 2$  edges. Thus  $\mathcal{R}(\mathcal{L}_\eta)$  has  $5\eta - 2$  vertices and  $3(3\eta - 2)$  edges. Therefore  $\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))$  has  $3(3\eta - 2)$  vertices and  $33\eta - 38$  edges.

#### *The edge partition of $\mathcal{L}(\mathcal{R}(\mathcal{L}_\eta))$*

Degree of end vertices	No. of edges
(4, 4)	6
(4, 6)	12
(4, 8)	8
(6, 6)	$9\eta - 20$
(6, 8)	16
(6, 10)	$18\eta - 48$
(8, 10)	8
(10, 10)	$(6\eta - 20)$

FIGURE 6.  $\mathcal{R}(\mathcal{L}_4)$ FIGURE 7.  $\mathcal{L}(\mathcal{R}(\mathcal{L}_4))$ 

The proof is similar as above.

□

## 6. CONCLUSION

To sum up we have calculated certain topological indices like Atom bond Connectivity index, Randic index, sum connectivity index, Geometric Arithmetic index, Sombor Index, First Zagreb index, Second Zagreb index, First Gourava index, Second Gourava index of Line graph of  $\mathcal{R}(\mathcal{G})$  in this paper. Computation of topological indices of chemical structures is still a scope open for future studies which help in QSPR or QSAR analysis.

## REFERENCES

1. Gutman, I. (2021). Geometric approach to degree-based topological indices: Sombor indices. MATCH Commun. Math. Comput. Chem, 86(1), 11-16.
2. Estrada, E., Torres, L., Rodriguez, L., & Gutman, I. (1998). An atom-bond connectivity index: modelling the enthalpy of formation of alkanes.
3. Randic, M. (1975). Characterization of molecular branching. Journal of the American Chemical Society, 97(23), 6609-6615.

4. Zhou, B., and Trinajstić, N. (2009). On a novel connectivity index. *Journal of mathematical chemistry*, 46, 1252-1270.
5. Vukičević, D. and Furtula, B. (2009). Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges. *Journal of mathematical chemistry*, 46, 1369-1376.
6. Gutman, I. and Trinajstić, N. (1972). Graph theory and molecular orbitals. Total  $\phi$ -electron energy of alternant hydrocarbons. *Chemical physics letters*, 17(4), 535-538.
7. Kulli, V. R. (2017). The Gourava indices and coindices of graphs. *Annals of Pure and Applied Mathematics*, 14(1), 33-38.
8. Adnan, M., Bokhary, S. A. U. H., Abbas, G., & Iqbal, T. (2022). Degree-Based Topological Indices and QSPR Analysis of Antituberculosis Drugs. *Journal of Chemistry*, 2022(1), 5748626.
9. Bokhary, S. A. U. H., Adnan, Siddiqui, M. K., & Cancan, M. (2022). On topological indices and QSPR analysis of drugs used for the treatment of breast cancer. *Polycyclic Aromatic Compounds*, 42(9), 6233-6253.
10. Mondal, S., De, N., & Pal, A. (2022). Topological indices of some chemical structures applied for the treatment of COVID-19 patients. *Polycyclic Aromatic Compounds*, 42(4), 1220-1234.
11. Khalaf, A. J. M., Hanif, M. F., Siddiqui, M. K., & Farahani, M. R. (2020). On degree based topological indices of bridge graphs. *Journal of Discrete Mathematical Sciences and Cryptography*, 23(6), 1139-1156.
12. Nasir, S., Farooq, F. B., & Parveen, S. (2022). Topological indices of novel drugs used in blood cancer treatment and its QSPR modeling. *AIMS Mathematics*, 7(7), 11829-11850.

<sup>1</sup> DEPARTMENT OF MATHEMATICS, ALL SAINTS' COLLEGE, THIRUVANATHAPURAM & NSS HINDU COLLEGE, CHANGANASSERY, KOTTAYAM.

*Email address:* [sebinayesu@gmail.com](mailto:sebinayesu@gmail.com)

<sup>2</sup> DEPARTMENT OF MATHEMATICS, NSS HINDU COLLEGE, CHANGANASSERY, KOTTAYAM

*Email address:* [satheeshkizhakkemadom@gmail.com](mailto:satheeshkizhakkemadom@gmail.com)