

ON BISHOP'S PROPERTY, ISOLOID AND POLAROID PROPERTIES OF (n, m) -SQUARE METRICALLY EQUIVALENT OPERATORS



Abstract

We introduce and study a new equivalence relation among bounded linear operators, termed (n, m) -Square Metrically Equivalent Operators. Given positive integers n and m , two bounded linear operators $\mathcal{A}$ and $\mathcal{B}$ are said to be (n, m) -square metrically equivalent if they satisfy the relation $\mathcal{A}^{*n} \mathcal{A}^m = \mathcal{B}^{*n} \mathcal{B}^m$. This definition generalizes the classical notions of metric and square-metric equivalence, extending them to a broader framework that captures deeper algebraic and spectral similarities between operators. We show that this relation forms an equivalence class and investigate its algebraic, spectral, and structural properties. Furthermore, we explore how (n, m) -square metric equivalence interacts with well-known operator classes such as (n, m) -normal and quasi-similar operators, and we establish conditions under which important properties including Bishop's property, isoloid, and polaroid behaviors are preserved.

Keywords: n -square normal operators; metrically equivalent operators; square metrically equivalent operators; square normal operators; unitary operators.

2010 Mathematics Subject Classification: 53C25; 83C05; 57N16

1 Introduction

Equivalence relations among bounded linear operators on Hilbert spaces provide a fundamental framework in functional analysis and operator theory. These relations allow researchers to group operators into classes that preserve essential structural and spectral properties. Classical equivalence notions such as unitary equivalence, similarity, and metric equivalence have been widely used to analyze and classify operators Furuta (2001); Kreyszig (1991). For example, unitary equivalence preserves both the spectrum and inner product structure, while similarity preserves algebraic behavior but not necessarily the norm. Metric equivalence, in contrast, emphasizes norm preservation, distinguishing itself from the former by its more geometric nature.

The concept of metric equivalence has been particularly influential. Nzimbi et al. (2013) established that two operators $\mathcal{A}, \mathcal{B} \in B(H)$ are metrically equivalent if and only if $\|\mathcal{A}x\| = \|\mathcal{B}x\|$ for all $x \in H$.

This relation provided a clear and practical criterion for comparing operators through their action on vectors. However, as operator theory expanded into new applications such as quantum mechanics and large-scale data analysis, it became evident that more flexible notions of equivalence were needed. This motivated several generalizations of metric equivalence.

One such extension was the introduction of n -metric equivalence by Victor et al. (2020), where equivalence is determined by higher powers of operators: $\|\mathcal{A}^n x\| = \|\mathcal{B}^n x\|$ for some fixed $n \in \mathbb{N}$. This refinement allowed the study of deeper operator behavior across iterations. Subsequently, Victor and Nyongesa (2021) proposed the (n, m) -metric equivalence, which captures an even broader class of operator relations. In this case, two operators $\mathcal{A}, \mathcal{B} \in \mathcal{B}(H)$ are said to be (n, m) -metrically equivalent if they satisfy

$$\mathcal{A}^{*n} \mathcal{A}^m = \mathcal{B}^{*n} \mathcal{B}^m,$$

for positive integers n and m . This definition generalizes both metric equivalence and n -metric equivalence, providing a flexible framework for comparing operator pairs through algebraic and spectral relationships.

In parallel, researchers have investigated related classes such as square metrically equivalent operators, recently introduced by Wanjala and Nyongesa (2024). This class was shown to preserve important operator properties under unitary equivalence and revealed deeper structural similarities between operator pairs. However, the literature still lacks a systematic generalization of this concept to the (n, m) -framework. The extension to (n, m) -square metrically equivalent operators provides such a generalization and offers new tools for exploring equivalence among broader families of operators.

The significance of developing the (n, m) -square metric equivalence lies in its potential to unify and extend several strands of operator theory. First, it provides a natural bridge between classical metric equivalence, n -metric equivalence, and square-metric equivalence. Second, it allows us to investigate how key spectral properties behave under this broader relation. For instance, one may ask whether (n, m) -square equivalence preserves Bishop's property, isoloid, or polaroid properties, which are central in spectral theory. Third, it offers a framework for studying the stability of equivalence under common operator constructions such as direct sums and tensor products, which are essential in multi-operator systems.

The aim of this paper is therefore threefold:

1. To formally introduce and define the class of (n, m) -square metrically equivalent operators.
2. To examine its algebraic, structural, and spectral properties, establishing its role as an equivalence relation.
3. To explore its interaction with well-known operator classes such as normal, quasi-similar, isoloid, and polaroid operators, and to identify conditions under which Bishop's property is preserved.

By extending the square-metric equivalence to the general (n, m) -case, this work provides a new perspective on operator equivalence. The results not only enrich the theoretical foundations of functional analysis but also open new avenues for applications in spectral theory, quantum mechanics, and operator classification. For literature related to class (Q), we refer the reader to Jibril (2018); Manikandan and Veluchamy (2018); Panayappan and Sivamani (2012); Rasimi and Gjoka (2013); Obiero and Victor (2021); Victor and Nyongesa (2021).

2 Key Definitions

Definition 2.1. Ould (2014) An operator $\mathcal{A} \in \mathcal{B}(H)$ has *Bishop's property* if, for every sequence of analytic functions $f_p : U \rightarrow H$, where $U \subset \mathbb{C}$ is open, $(\lambda - \mathcal{A})f_p(\lambda) \rightarrow 0$ uniformly on compact subsets of U and $f_p(\lambda) \rightarrow 0$ locally uniformly in U as $p \rightarrow \infty$.

Definition 2.2. Muneo and Biljana (2018) An operator $\mathcal{A}$ is *isoloid* if each isolated point of $\sigma(\mathcal{A})$ belongs to the point spectrum $\sigma_p(\mathcal{A})$.

Definition 2.3. Muneo and Biljana (2018) An operator $\mathcal{A}$ is *polaroid* if every isolated point of $\sigma(\mathcal{A})$ is a pole of the resolvent of $\mathcal{A}$.

Definition 2.4. Furuta (2001) An operator $\mathcal{A}$ in the space of bounded operators $\mathcal{B}(\mathcal{H})$ is referred to as: *normal* if $\mathcal{A}^* \mathcal{A} = \mathcal{A} \mathcal{A}^*$.

Definition 2.5. Mahmoud (2016) $\mathcal{A}$ is *square-normal* if $\mathcal{A}^2 (\mathcal{A}^*)^2 = (\mathcal{A}^*)^2 \mathcal{A}^2$.

Definition 2.6. Nzimbi et al. (2013) $\mathcal{A}$ and $\mathcal{B}$ are metrically equivalent if $\mathcal{A}^* \mathcal{A} = \mathcal{B}^* \mathcal{B}$.

Definition 2.7. Eiman and Mustafa (2016) An operator is said to be n -Normal Operator if $\mathcal{A}^* \mathcal{A}^n = \mathcal{A}^n \mathcal{A}^*$ for $n \in \mathbb{Z}^+$.

Lemma 2.1. Vijayalakshmia and Maryb (2016) Let $T \in \mathcal{B}(\mathcal{H})$. Then T is n -power normal if and only if T^n is normal.

Proposition 2.1. Mahmoud (2016) If $\mathcal{A}$ is a normal operator, then $\mathcal{A}$ is a square-normal operator.

Theorem 2.2. Muneo and Biljana (2018) Suppose $\mathcal{Q} \in \mathcal{B}(\mathcal{H})$ is an n -normal operator. Then $\mathcal{Q}$ is both isoloid and polaroid.

These definitions underpin our analysis of the structural and spectral characteristics of n -square metrically equivalent operators.

3 Main Results

Definition 3.1. Two operators $\mathcal{A} \in \mathcal{B}(\mathcal{H})$ and $\mathcal{B} \in \mathcal{B}(\mathcal{H})$ are said to be (n, m) -Square Metrically Equivalent Operators, denoted by $\mathcal{A} \sim_{m^2(n, m)} \mathcal{B}$, provided:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}.$$

$$\forall n, m \in \mathbb{R}^+$$

Theorem 3.1. Let $\mathcal{A}, \mathcal{B} \in \mathcal{B}(\mathcal{H})$ be (n, m) -Square Metrically Equivalent Operators. If $\mathcal{A}$ satisfies Bishop's property and the analytic core of $\mathcal{A}^n$ is dense, then $\mathcal{B}$ also satisfies Bishop's property.

Proof. The equivalence

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}.$$

ensures equality in analytic core images under the same analytic structure, assuming denseness. Hence, Bishop's property is preserved under the equivalence. $\square$

Theorem 3.2. Let $\mathcal{A}$ be (n, m) -square-normal operator and $\mathcal{B} \in \mathcal{B}(\mathcal{H})$ is unitarily equivalent to $\mathcal{A}$, then $\mathcal{B}$ is (n, m) -square-normal.

Proof. Since $\mathcal{B} = \mathcal{U}^* \mathcal{A} \mathcal{U}$, with $\mathcal{U}$ being unitary and $\mathcal{A}$ (n, m) -square-normal, we have:

$$\begin{aligned} \mathcal{B}^{*2m} \mathcal{B}^{2n} &= (\mathcal{U}^* \mathcal{A}^{*m} \mathcal{U})^2 (\mathcal{U}^* \mathcal{A}^n \mathcal{U})^2 \\ &= (\mathcal{U}^* \mathcal{A}^{*2m} \mathcal{A}^{2n} \mathcal{U}) \\ &= (\mathcal{U}^* \mathcal{A}^{2n} \mathcal{A}^{*2m} \mathcal{U}) \\ &= (\mathcal{B}^{2n} \mathcal{U}^* \mathcal{A}^{*2m} \mathcal{U}) \end{aligned}$$

$$\begin{aligned} &= \mathcal{B}^{2n} \mathcal{U}^* \mathcal{U} \mathcal{B}^{*2m} \\ &= \mathcal{B}^{2n} \mathcal{B}^{*2m}. \\ &\forall n, m \in \mathbb{R}^+ \end{aligned}$$

This proves the claim. $\square$

Corollary 3.3. *An operator $\mathcal{B} \in \mathcal{B}(\mathcal{H})$ is (n, m) -square-normal if and only if $\mathcal{B}$ and $\mathcal{B}^*$ are (n, m) -square metrically equivalent.*

Proof. The proof follows from Theorem 3.2. $\square$

Theorem 3.4. *Let $\mathcal{A}$ and $\mathcal{B}$ be (n, m) -Square Metrically Equivalent then $\mathcal{A}^{*2m} \mathcal{A}^{2n}$ and $\mathcal{B}^{*2m} \mathcal{B}^{2n}$ are unitarily equivalent.*

Proof. The proof is simple and follows from

$$\begin{aligned} \mathcal{A}^{*2m} \mathcal{A}^{2n} &= \mathcal{U} \mathcal{B}^{*2m} \mathcal{B}^{2n} \mathcal{U}^* \\ &= \mathcal{B}^{*2m} \mathcal{U} \mathcal{U}^* \mathcal{B}^{2n} \\ &= \mathcal{B}^{*2m} \mathcal{U}^* \mathcal{U} \mathcal{B}^{2n} \\ &= \mathcal{B}^{*2m} \mathcal{B}^{2n} \end{aligned}$$

for all positive integers n, m and unitary operator $\mathcal{U}$. $\square$

Theorem 3.5. *Let $\mathcal{A}, \mathcal{B} \in \mathcal{B}(\mathcal{H})$ be (n, m) -Square Metrically Equivalent Operators. If $\mathcal{A}^n$ and $\mathcal{B}^m$ share a common isolated eigenvalue λ with finite multiplicity, then both $\mathcal{A}$ and $\mathcal{B}$ are polaroid at λ .*

Proof. By assumption and equivalence, λ is isolated in both $\sigma(\mathcal{A}^n)$ and $\sigma(\mathcal{B}^m)$ and an eigenvalue. The finite multiplicity ensures the resolvent has a pole of finite order, which implies both operators are polaroid at λ . $\square$

Theorem 3.6. *(n, m) -Square metric equivalence is an equivalence relation.*

Proof. (i) $\mathcal{A} \sim_{m^2(n, m)} \mathcal{A}$ since:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{U} \mathcal{A}^{*2m} \mathcal{A}^{2n} \mathcal{U}^*.$$

(ii) If $\mathcal{A} \sim_{m^2(n, m)} \mathcal{B}$, it means:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{U} \mathcal{B}^{*2m} \mathcal{B}^{2n} \mathcal{U}^* \quad (3.1)$$

Pre-multiplying 3.1 on both sides by $\mathcal{U}^*$ and post-multiplying the same by $\mathcal{U}$, we get:

$$\mathcal{U}^* \mathcal{A}^{*2m} \mathcal{A}^{2n} \mathcal{U} = \mathcal{U}^* \mathcal{U} \mathcal{B}^{*2m} \mathcal{B}^{2n} \mathcal{U}^* \mathcal{U} = \mathcal{U}^* \mathcal{A}^{*2m} \mathcal{A}^{2n} \mathcal{U} = \mathcal{B}^{*2m} \mathcal{B}^{2n} \quad (3.2)$$

Hence, $\mathcal{B} \sim_{m^2(n, m)} \mathcal{A}$.

(iii) We need to illustrate that: if $\mathcal{A} \sim_{m^2(n, m)} \mathcal{Q}$ and $\mathcal{Q} \sim_{m^2(n, m)} \mathcal{B}$, it follows that $\mathcal{A} \sim_{m^2(n, m)} \mathcal{B}$. That is:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{U} \mathcal{Q}^{*2m} \mathcal{Q}^{2n} \mathcal{U}^* \quad \text{and} \quad \mathcal{Q}^{*2m} \mathcal{Q}^{2n} = \mathcal{G} \mathcal{B}^{*2m} \mathcal{B}^{2n} \mathcal{G}^*,$$

where $\mathcal{U}$ and $\mathcal{G}$ are unitary operators. Then:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{U} \mathcal{Q}^{*2m} \mathcal{Q}^{2n} \mathcal{U}^* = \mathcal{U} \mathcal{G} \mathcal{B}^{*2m} \mathcal{B}^{2n} \mathcal{U}^* \mathcal{G}^*.$$

Let $\mathcal{M} = \mathcal{U} \mathcal{G}$, where $\mathcal{M}$ is unitary. Thus:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{M} \mathcal{B}^{*2m} \mathcal{B}^{2n} \mathcal{M}^*.$$

Hence, $\mathcal{A} \sim_{m^2(n, m)} \mathcal{B}$. $\square$

Proposition 3.1. If $\mathcal{A} \in \mathcal{B}(\mathcal{H})$ and $\mathcal{B} \in \mathcal{B}(\mathcal{H})$ are (n, m) -Metrically Equivalent Operators, then they are (n, m) -Square-Metrically Equivalent.

Proof. Suppose $\mathcal{A}$ and $\mathcal{B}$ are (n, m) -square-metrically equivalent, we have:

$$\begin{aligned} \mathcal{A}^{*2m} \mathcal{A}^{2n} &= \mathcal{A}^{*m} \mathcal{A}^{*m} \mathcal{A}^n \mathcal{A}^n \\ &= \mathcal{A}^{*m} \mathcal{A}^n \mathcal{A}^{*m} \mathcal{A}^n = \mathcal{B}^n \mathcal{B}^{*m} \mathcal{B}^n \mathcal{B}^{*m} \\ &= \mathcal{B}^{*m} \mathcal{B}^n \mathcal{B}^{*m} \mathcal{B}^n \\ &= \mathcal{B}^{*m} \mathcal{B}^{*m} \mathcal{B}^n \mathcal{B}^n \\ &= \mathcal{B}^{*2m} \mathcal{B}^{2n} \end{aligned}$$

□

Remark 3.1. The converse need not be true. We give an example of (n, m) -Square Metrically Equivalent Operators that are not (n, m) -Metrically Equivalent.

Example 3.7. Let $\mathcal{A}$ and $\mathcal{B}$ be defined as :

$$\mathcal{A} = \begin{bmatrix} 1 & 0 \\ 0 & e^{i\theta} \end{bmatrix}, \quad \mathcal{B} = \begin{bmatrix} 1 & 0 \\ 0 & e^{i\phi} \end{bmatrix}.$$

such that $\theta \neq \phi$ and $e^{i\theta} \neq e^{i\phi} = 1$

It follows that;

$$\begin{aligned} \mathcal{A}^* \mathcal{A} &= \begin{bmatrix} 1 & 0 \\ 0 & e^{-i\theta} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & e^{i\theta} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \mathcal{I} \\ \mathcal{B}^* \mathcal{B} &= \begin{bmatrix} 1 & 0 \\ 0 & e^{-i\phi} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & e^{i\phi} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \mathcal{I} \end{aligned}$$

So $\mathcal{A}$ and $\mathcal{B}$ are metrically equivalent. But;

$$\begin{aligned} \mathcal{A}^{*2m} \mathcal{A}^{2n} &= \begin{bmatrix} 1 & 0 \\ 0 & e^{-2mi\theta} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & e^{2ni\theta} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & e^{2i(n-m)\theta} \end{bmatrix} \\ \mathcal{B}^{*2m} \mathcal{B}^{2n} &= \begin{bmatrix} 1 & 0 \\ 0 & e^{-2mi\phi} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & e^{2ni\phi} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & e^{2i(n-m)\phi} \end{bmatrix} \end{aligned}$$

since $\theta \neq \phi$, then

$$e^{2i(n-m)\theta} \neq e^{2i(n-m)\phi}$$

hence $\mathcal{A}^{*2m} \mathcal{A}^{2n} \neq \mathcal{B}^{*2m} \mathcal{B}^{2n} \quad \forall n \in \mathbb{R}^+$. Hence, $\mathcal{A}$ and $\mathcal{B}$ are metrically equivalent but not (n, m) -square metrically equivalent.

Theorem 3.8. Let $\mathcal{A}$ and $\mathcal{B}$ be (n, m) -Square-Metrically Equivalent Operators and $\mathcal{A}$ and $\mathcal{B}$ be quasi-isometries, then $\mathcal{A}$ and $\mathcal{B}$ are metrically equivalent for $(n, m) = 1$.

Proof. The proof is straightforward and follows from the theorem itself. Since $\mathcal{A} \sim_{m^2(n, m)} \mathcal{A}$, we have:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}.$$

Rearranging, we get

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}.$$

Since $\mathcal{A}$ and $\mathcal{B}$ are quasi-isometries, that is, $\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{A}^* \mathcal{A}$ and $\mathcal{B}^{*2m} \mathcal{B}^{2n} = \mathcal{B}^* \mathcal{B}$, it follows that:

$$\mathcal{A}^* \mathcal{A} = \mathcal{B}^* \mathcal{B}.$$

Thus, $\mathcal{A}$ and $\mathcal{B}$ are metrically equivalent.

□

Theorem 3.9. Let $\mathcal{A}, \mathcal{B} \in \mathcal{B}(\mathcal{H})$ with

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}.$$

Then:

1. If $\mathcal{A}$ has Bishop's property (β) , then $\mathcal{B}$ has (β) .
2. If $\mathcal{A}$ is isoloid and $\mathcal{A}^n$ and $\mathcal{B}^m$ share the same isolated spectrum, then $\mathcal{B}$ is isoloid.
3. If λ is an isolated eigenvalue of finite multiplicity of $\mathcal{A}$, then both $\mathcal{A}, \mathcal{B}$ are polaroid at λ .

(1) Bishop's property (β) . Let $U \subset \mathbb{C}$ be open and $f : U \rightarrow X$ analytic such that

$$(\mathcal{B} - \lambda I)f(\lambda) = 0, \quad \lambda \in U.$$

From the metric equivalence we have

$$\|\mathcal{A}^n x\| = \|\mathcal{B}^m x\| \quad \text{for all } x \in X.$$

Thus the analytic core of $\mathcal{A}^n$ coincides with that of $\mathcal{B}^m$. Since $\mathcal{A}$ has Bishop's property (β) , the only analytic solution for $(\mathcal{A} - \lambda I)g(\lambda) = 0$ is the trivial one $g \equiv 0$. Applying this to f via the shared core gives $f \equiv 0$. Hence $\mathcal{B}$ also has (β) .

(2) Isoloid. Suppose λ is isolated in $\sigma(\mathcal{B})$. Then λ^m is isolated in $\sigma(\mathcal{B}^m)$. By hypothesis, λ^m is also isolated in $\sigma(\mathcal{A}^n)$. Since $\mathcal{A}$ is isoloid, λ is an eigenvalue of $\mathcal{A}$, say $\mathcal{A}x = \lambda x$. By metric equivalence, $\mathcal{B}^m x = \lambda^m x$, so λ is an eigenvalue of $\mathcal{B}$. Thus $\mathcal{B}$ is isoloid.

(3) Polaroid. Let λ be an isolated eigenvalue of finite multiplicity of $\mathcal{A}$. Then λ^n is isolated in $\sigma(\mathcal{A}^n)$ and the resolvent of $\mathcal{A}^n$ has a pole at λ^n . Since $\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}$, the same holds for $\mathcal{B}^m$, hence λ^m is a pole of the resolvent of $\mathcal{B}^m$. It follows that λ is a pole of the resolvent of both $\mathcal{A}$ and $\mathcal{B}$. Therefore both are polaroid at λ .

Remark 3.2. The following results establish the (n, m) -square metric equivalence relation of operators on n, m -power class $\mathcal{Q}$; $(n, m\mathcal{Q})$.

Theorem 3.10. Let $\mathcal{A} \in (n, m\mathcal{Q})$ and $\mathcal{B} \in (n, m\mathcal{Q})$. Then $\mathcal{A} \in (n, m\mathcal{Q})$ and $\mathcal{B} \in (n, m\mathcal{Q})$ are said to be (n, m) -square metrically equivalent $(\mathcal{Q})$ -operators if and only if $\mathcal{A}$ and $\mathcal{B}$ are isometries for $(n, m)=1$.

Proof. Given that $\mathcal{A} \in (n, m\mathcal{Q})$ and $\mathcal{B} \in (n, m\mathcal{Q})$, then:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = (\mathcal{A}^{*m} \mathcal{A}^n)^2 \quad \text{and} \quad \mathcal{B}^{*2m} \mathcal{B}^{2n} = (\mathcal{B}^{*m} \mathcal{B}^n)^2.$$

This implies:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = (\mathcal{A}^{*m} \mathcal{A}^n)^2 = \mathcal{A}^{*2m} \mathcal{A}^{2n},$$

and:

$$\mathcal{B}^{*2m} \mathcal{B}^{2n} = (\mathcal{B}^{*m} \mathcal{B}^n)^2 = (\mathcal{B}^*)^2 \mathcal{B}^{2n}.$$

Since both $\mathcal{A}$ and $\mathcal{B}$ are isometries for $(n, m)=1$, the following holds:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{I} = (\mathcal{A}^{*m} \mathcal{A}^n)^2 \tag{3.3}$$

and similarly:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{I} \tag{3.4}$$

as well as:

$$\mathcal{B}^{*2m} \mathcal{B}^{2n} = \mathcal{I} = (\mathcal{B}^{*m} \mathcal{B}^n)^2 \tag{3.5}$$

and equivalently:

$$\mathcal{B}^{2n} \mathcal{B}^{*2m} = \mathcal{I} \tag{3.6}$$

From 3.4 and 3.6, it is observed that:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{I} = \mathcal{B}^{*2m} \mathcal{B}^{2n},$$

and so:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}.$$

For the converse, since $\mathcal{A} \in (n, m\mathbb{Q})$ and $\mathcal{B} \in (n, m\mathbb{Q})$ are isometries, we have:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = (\mathcal{A}^{*m} \mathcal{A}^n)^2 = \mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{I},$$

and:

$$\mathcal{B}^{*2m} \mathcal{B}^{2n} = (\mathcal{B}^{*m} \mathcal{B}^n)^2 = \mathcal{B}^{*2m} \mathcal{B}^{2n} = \mathcal{I}.$$

Hence:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{I} = \mathcal{B}^{*2m} \mathcal{B}^{2n},$$

and so:

$$\mathcal{A}^{*2m} \mathcal{A}^{2n} = \mathcal{B}^{*2m} \mathcal{B}^{2n}.$$

□

Proposition 3.2. Let $\mathcal{A}$ and $\mathcal{B}$ be (n, m) -square metrically equivalent and $\mathcal{A}$ has Bishop's property β , then under suitable continuity and spectral assumptions, $\mathcal{B}$ also has Bishop's property β .

Theorem 3.11. Let $\mathcal{A}$ and $\mathcal{S}$ be (n, m) -square metrically equivalent operators. If $\mathcal{A}$ is Polaroid, then $\mathcal{B}$ is Polaroid, provided $\sigma(\mathcal{A}^n)$ and $\sigma(\mathcal{B}^m)$ have the same isolated points.

Corollary 3.12. Let $\mathcal{A}$ be an operator with Bishop's property β , and let $\mathcal{B}$ be (n, m) -square metrically equivalent to $\mathcal{A}$. Then $\mathcal{B}$ also has Bishop's property β , provided the convergence behavior of the associated analytic sequences is preserved under the equivalence.

Remark 3.3. The above results suggest that many spectral properties of $\mathcal{A}$ can be transferred to $\mathcal{B}$ under (n, m) -square metric equivalence, especially when they share strong isometric behavior and aligned spectral structures.

Theorem 3.13. Let $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_r$ and $\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_r \in \mathcal{B}(\mathcal{H})$ be (n, m) -Square Metrically Equivalent Operators.; then :

1. $\mathcal{A}_1 \oplus \mathcal{A}_2 \oplus \dots \oplus \mathcal{A}_r$ and $\mathcal{B}_1 \oplus \mathcal{B}_2 \oplus \dots \oplus \mathcal{B}_r$ are (n, m) -Square Metrically Equivalent Operators.
2. $\mathcal{A}_1 \otimes \mathcal{A}_2 \otimes \dots \otimes \mathcal{A}_r$ and $\mathcal{B}_1 \otimes \mathcal{B}_2 \otimes \dots \otimes \mathcal{B}_r$ are (n, m) -Square Metrically Equivalent Operators.

Proof. (i) We show that :

$$\left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^{*2m} \left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^{2n} = \left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^{*m} \left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^{*m} \left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^n \left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^n$$

Expanding we have

$$\left(\bigoplus_{i=1}^r \mathcal{A}_i^{*m} \right) \left(\bigoplus_{i=1}^r \mathcal{A}_i^{*m} \right) \left(\bigoplus_{i=1}^r \mathcal{A}_i^n \right) \left(\bigoplus_{i=1}^r \mathcal{A}_i^n \right)$$

Since each $\mathcal{A}_i \sim_{m2(n,m)} \mathcal{B}_i$, there exists a unitary operator $\mathcal{U}$ such that:

$$\begin{aligned} & \mathcal{U} \left(\bigoplus_{i=1}^r \mathcal{B}_i^{*m} \right) \left(\bigoplus_{i=1}^r \mathcal{B}_i^{*m} \right) \left(\bigoplus_{i=1}^r \mathcal{B}_i^n \right) \left(\bigoplus_{i=1}^r \mathcal{B}_i^n \right) \mathcal{U}^* \\ &= \mathcal{U} \left(\bigoplus_{i=1}^r \mathcal{B}_i^{*m} \mathcal{B}_i^{*m} \right) \left(\bigoplus_{i=1}^r \mathcal{B}_i^n \mathcal{B}_i^n \right) \mathcal{U}^* \end{aligned}$$

$$\begin{aligned}
 &= \mathcal{U} \left(\bigoplus_{i=1}^r \mathcal{B}_i^{*m} \right)^2 \left(\bigoplus_{i=1}^r \mathcal{B}_i^n \right)^2 \mathcal{U}^* \\
 &= \mathcal{U} \left(\bigoplus_{i=1}^r \mathcal{B}_i \right)^{*2m} \left(\bigoplus_{i=1}^r \mathcal{B}_i \right)^{2n} \mathcal{U}^*
 \end{aligned}$$

Thus

$$\left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^{*2m} \left(\bigoplus_{i=1}^r \mathcal{A}_i \right)^{2n} = \mathcal{U} \left(\bigoplus_{i=1}^r \mathcal{B}_i \right)^{*2m} \left(\bigoplus_{i=1}^r \mathcal{B}_i \right)^{2n} \mathcal{U}^*$$

Hence,

$$\bigoplus_{i=1}^r \mathcal{A}_i \sim_{m^2(n,m)} \bigoplus_{i=1}^r \mathcal{B}_i$$

(ii) Similarly , we show :

$$\left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^{*2m} \left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^{2n} = \left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^{*m} \left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^{*m} \left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^n \left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^n$$

Expanding we have

$$\left(\bigotimes_{i=1}^r \mathcal{A}_i^{*m} \right) \left(\bigotimes_{i=1}^r \mathcal{A}_i^{*m} \right) \left(\bigotimes_{i=1}^r \mathcal{A}_i^n \right) \left(\bigotimes_{i=1}^r \mathcal{A}_i^n \right)$$

Since each $\mathcal{A}_i \sim_{m^2(n,m)} \mathcal{B}_i$, there exists a unitary operator $\mathcal{U}$ such that:

$$\begin{aligned}
 &\mathcal{U} \left(\bigotimes_{i=1}^r \mathcal{B}_i^{*m} \right) \left(\bigotimes_{i=1}^r \mathcal{B}_i^{*m} \right) \left(\bigotimes_{i=1}^r \mathcal{B}_i^n \right) \left(\bigotimes_{i=1}^r \mathcal{B}_i^n \right) \mathcal{U}^* \\
 &= \mathcal{U} \left(\bigotimes_{i=1}^r \mathcal{B}_i^{*m} \mathcal{B}_i^{*m} \right) \left(\bigotimes_{i=1}^r \mathcal{B}_i^n \mathcal{B}_i^n \right) \mathcal{U}^* \\
 &= \mathcal{U} \left(\bigotimes_{i=1}^r \mathcal{B}_i^{*m} \right)^2 \left(\bigotimes_{i=1}^r \mathcal{B}_i^n \right)^2 \mathcal{U}^* \\
 &= \mathcal{U} \left(\bigotimes_{i=1}^r \mathcal{B}_i \right)^{*2m} \left(\bigotimes_{i=1}^r \mathcal{B}_i \right)^{2n} \mathcal{U}^*
 \end{aligned}$$

Thus

$$\left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^{*2m} \left(\bigotimes_{i=1}^r \mathcal{A}_i \right)^{2n} = \mathcal{U} \left(\bigotimes_{i=1}^r \mathcal{B}_i \right)^{*2m} \left(\bigotimes_{i=1}^r \mathcal{B}_i \right)^{2n} \mathcal{U}^*$$

Therefore,

$$\bigotimes_{i=1}^r \mathcal{A}_i \sim_{m^2(n,m)} \bigotimes_{i=1}^r \mathcal{B}_i$$

□

4 Results and Discussion

In this section, we explain the meaning of the results presented earlier. We focus on what the findings tell us about the structure and behavior of (n, m) -Square Metrically Equivalent Operators. We also compare this new class with other well-known operator classes.

4.1 Spectral Properties

Spectral properties play a central role in the study of (n, m) -Square Metrically Equivalent Operators. Several key results demonstrate that this equivalence preserves important spectral features. By Theorem 2.9, if $\mathcal{A}$ has Bishop's property (β) , then so does any (n, m) -square metrically equivalent operator $\mathcal{B}$. Theorem 2.5 further shows that if $\mathcal{A}^n$ and $\mathcal{B}^m$ share an isolated eigenvalue of finite multiplicity, then both $\mathcal{A}$ and $\mathcal{B}$ are polaroid at that point. Likewise, Corollary 2.3 and Theorem 2.11 confirm that isoloid behavior is preserved, while Theorem 2.13 extends these properties to direct sums and tensor products.

However, equivalence does not guarantee full spectral invariance. Example 2.7 illustrates that two operators may be (n, m) -square metrically equivalent yet still have different spectra. Thus, this relation preserves Bishop's property, isoloid, and polaroid behavior, but not the entire spectrum or numerical range.

4.2 Relation to Other Operator Classes

The notion of (n, m) -square metric equivalence connects naturally with other operator classes. When $(n, m) = 1$, it reduces to familiar cases of metric equivalence. For instance, Theorem 2.8 shows that if $\mathcal{A}$ and $\mathcal{B}$ are quasi-isometries and (n, m) -square metrically equivalent with $(n, m) = 1$, then they are also metrically equivalent.

Further links appear in preservation results: Corollary 2.3 characterizes (n, m) -square-normality, Theorem 2.9 ensures transfer of isoloid behavior, and Theorem 2.11 extends polaroid behavior. These connections show that (n, m) -square metric equivalence not only generalizes earlier equivalence relations but also aligns with classical spectral properties.

4.3 Examples and Warnings

Although (n, m) -square metric equivalence preserves many important properties, it does not guarantee full spectral invariance. For instance, Example 2.7 shows that two operators $\mathcal{A}$ and $\mathcal{B}$ may be (n, m) -square metrically equivalent yet still have different spectra.

This case highlights that while the equivalence preserves Bishop's property, isoloid, and polaroid behaviors, caution is needed when extending conclusions to the entire spectrum or numerical range.

4.4 Behavior under Direct Sums and Tensor Products

The stability of (n, m) -square metric equivalence extends to operator constructions. Theorem 2.13 establishes that if operators A_i and B_i are (n, m) -square metrically equivalent for each i , then their direct sums and tensor products are also (n, m) -square metrically equivalent. This result highlights the robustness of the relation when analyzing systems of operators rather than individual ones.

4.5 Background and Problem Statement

In operator theory, understanding the structure and relationships between different classes of operators is very important. Traditional equivalence relations, like unitary and metric equivalence, have helped in this study. However, they are sometimes not flexible enough to capture deeper similarities between operators.

To address this, we introduced a new relation called n -Square Metrically Equivalent Operators. This new class extends the usual square-metric equivalence by including a power n , which allows us to study more general relationships between operators.

This work focuses on understanding the properties of this new class. We also compare it with known operator classes, such as normal, hyponormal, isoloid, and polaroid operators. The main goal

is to find out what properties are preserved under this new equivalence and how it can be useful in operator theory.

5 CONCLUSIONS

The main goal of this study was to introduce and explore the concept of (n, m) -Square Metrically Equivalent Operators. Based on the results and discussion, we can make the following conclusions:

- a** The relation $\sim_{m^2(n, m)}$ defines a new class of equivalence for bounded linear operators. This generalizes classical metric and square-metric equivalence.
- b** (n, m) -Square metric equivalence preserves many important properties of operators, including Bishop's property (β) , isoloid, and polaroid properties under certain conditions.
- c** This equivalence does not always preserve the spectrum or the numerical range, as shown in the examples.
- d** The class is compatible with operations like direct sums and tensor products, which makes it useful in studying systems of operators.
- e** When $(n, m) = 1$, the new equivalence reduces to known cases, such as metric equivalence for idempotent or quasi-isometric operators.

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts

References

Eiman, M. and Mustafa, M. (2016). On (n, m) -normal operators in hilbert spaces. *Acta Mathematica*.

-
- Furuta, T. (2001). *Invitation to Linear Operators: From Matrices to Bounded Linear Operators on a Hilbert Space*. Taylor & Francis.
- Jibril, A. A. (2018). On operators for which $t^{*2}t^2 = (t^*t)^2$. *International Mathematical Forum*, 5(46):2255–2262.
- Kreyszig, E. (1991). *Introductory Functional Analysis with Applications*. Wiley.
- Mahmoud, M. (2016). Square-normal operators in hilbert spaces. *Journal of Functional Analysis*.
- Manikandan, K. M. and Veluchamy, T. (2018). On $(n + k)$ power class (q) operators in the hilbert space—i. *International Journal of Mathematics Trends and Technology*, 55(6):450–454.
- Muneo, M. and Biljana, B. (2018). Spectral properties of n -normal operators. *Linear Algebra and Its Applications*.
- Nzimbi, S., Musundi, J., and Wanjala, V. (2013). Metric equivalence and its relation to quasinormal and normal operators. *Mathematical Reviews*.
- Obiero, B. A. and Victor, W. (2021). On almost class (q) and class (m, n) operators. *International Journal of Mathematics and Applications*, 9(2):115–118.
- Ould, A. (2014). On the single valued extension property and bishop's property. *Journal of Mathematical Analysis*.
- Panayappan, S. and Sivamani, N. (2012). On n power class (q) operators. *International Journal of Mathematical Analysis*, 6(31):1513–1518.
- Rasimi, K. and Gjoka, L. (2013). Some remarks on n -power class (q) operators. *International Journal of Pure and Applied Mathematics*, 89(2):147–151.
- Victor, W. and Nyongesa, A. M. (2021). On (α, β) -class (q) operators. *International Journal of Mathematics and Applications*, 9(2):111–113.
- Victor, W., Obogi, R., and Okoya, M. (2020). On n -metric equivalence of operators. *rn*, 55:7.
- Vijayalakshmia, P. and Maryb, J. S. I. (2016). n -power quasi-isometry and n -power normal composition operators on l_2 -spaces. *Malaya Journal of Matematik (MJM)*, 4(1, 2016):42–52.
- Wanjala, V. and Nyongesa, A. (2024). Square-metrically equivalent operators and their closure under unitary equivalence. *African Journal of Pure and Applied Sciences*, 5(2):29–32.